

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EcoRI
Catalog Number: R0101L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10119696
Expiration Date: 06/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM KPO4 (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0101S/L v2.0

EcoRI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0101LVIAL	EcoRI	10112498	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10114156	Pass	
B0101SVIAL	NEBuffer™ EcoRI/SspI	10110279	Pass	

Protein Purity Assay (SDS-PAGE) ECORI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue
CONT is 293 % pure as determined by 3D3 FAGE analysis using Coomassie blue
letection.
Blue-White Screening (Terminal Integrity) Pass
sample of pUC19 vector linearized with a 10-fold excess of EcoRI, religated and ransformed into an E. coli strain expressing the LacZ beta fragment gene results in
white colonies.
Exonuclease Activity (Radioactivity Release)
λ 50 μl reaction in EcoRl Buffer containing 1 μg of a mixture of single and
ouble-stranded [³H] E. coli DNA and a minimum of 200 units of EcoRI incubated for hours at 37°C releases <0.1% of the total radioactivity.
Ion-Specific DNase Activity (16 Hour) Pass
λ 50 μl reaction in EcoRI Buffer containing 1 μg of Lambda DNA and a minimum of 100
Inits of EcoRI incubated for 16 hours at 37°C results in a DNA pattern free of etectable nuclease degradation as determined by agarose gel electrophoresis.



R0101L / Lot: 10119696 Page 1 of 2

Assay Name/Specification	Lot # 10119696
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with EcoRI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRI.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

07 Sep 2021

Josh Hersey

Packaging Quality Control Inspector

07 Sep 2021

