

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	EcoRI
Catalog Number:	R0101M
Concentration:	100,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10135001
Expiration Date:	12/2023
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM KPO4 (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml BSA
Specification Version:	PS-R0101T/M v3.0

EcoRI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0101MVIAL	EcoRI	10135000	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10130600	Pass	
B0101SVIAL	NEBuffer™ EcoRI/SspI	10110279	Pass	

Assay Name/Specification	Lot # 10135001
Protein Purity Assay (SDS-PAGE) EcoRI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Blue-White Screening (Terminal Integrity)</b> A sample of pUC19 vector linearized with a 10-fold excess of EcoRI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with EcoRI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRI.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in 1X NEBuffer EcoRI/Sspl containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of EcoRI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10135001
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in 1X NEBuffer EcoRI/Sspl containing 1 µg of Lambda DNA and a minimum of 100 units of EcoRI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pefnilly

Penghua Zhang Production Scientist 08 Feb 2022

Michael Tonello Packaging Quality Control Inspector 08 Feb 2022

