

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: EcoRI
Catalog Number: R0101T
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10154599
Expiration Date: 05/2024
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM KPO4 (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0101T/M v3.0

EcoRI Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0101TVIAL	EcoRI	10150308	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10150372	Pass	
B0101SVIAL	NEBuffer™ EcoRI/SspI	10110279	Pass	

Assay Name/Specification	Lot # 10154599
Protein Purity Assay (SDS-PAGE)  EcoRI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	
Blue-White Screening (Terminal Integrity)	Pass
A sample of pUC19 vector linearized with a 10-fold excess of EcoRI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with EcoRI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRI.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in 1X NEBuffer EcoRl/Sspl containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 200 units of EcoRl incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R0101T / Lot: 10154599

Page 1 of 2

Assay Name/Specification	Lot # 10154599
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in 1X NEBuffer EcoRl/Sspl containing 1 µg of Lambda DNA and a minimum of 100 units of EcoRl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

06 Jun 2022

Erin Varney

Packaging Quality Control Inspector

06 Jun 2022

