

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HindIII

Catalog Number: R0104S

Concentration: 20,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10103327 Expiration Date: 01/2023 Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0104S/L v1.0

HindIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0104SVIAL	HindIII	10098120	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091457	Pass	
B6002SVIAL	NEBuffer™ r2.1	10103928	Pass	

Assay Name/Specification	Lot # 10103327
Protein Purity Assay (SDS-PAGE) HindIII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 60	Pass
Units of HindIII incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 DNA and a	Pass
minimum of 60 Units of HindIII incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 200 units of HindIII incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R0104S / Lot: 10103327

Page 1 of 2

Assay Name/Specification	Lot # 10103327
Ligation and Recutting (Terminal Integrity) After a 200-fold over-digestion of Lambda DNA with HindIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HindIII.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of HindIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang **Production Scientist**

06 Apr 2021

Michael Tonello

Packaging Quality Control Inspector

06 Apr 2021



R0104S / Lot: 10103327

Page 2 of 2