

## New England Biolabs Certificate of Analysis

**Product Name:** *FokI*  
**Catalog #:** *R0109S/L*  
**Concentration:** *5,000 units/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Lot #:** *0471501*  
**Assay Date:** *01/2015*  
**Expiration Date:** *01/2017*  
**Storage Temp:** *-20 °C*  
**Storage Conditions:** *50 mM NaCl, 10 mM Tris-HCl (pH 8.2), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.1% Tween-20, 200 µg/ml BSA*  
**Specification Version:** *PS-R0109S/L v1.0*  
**Effective Date:** *15 Jul 2013*

Assay Name/Specification (minimum release criteria)	Lot #0471501
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 5 units of FokI incubated for 4 hours at 37°C releases <0.5% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with FokI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with FokI.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
15 Jul 2013



Inspected by  
Anthony Francis  
30 Jan 2015

