

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | FokI  |
|------------------------|---|
| Catalog #:             | R0109S/L  |
| Concentration:         | 5,000 units/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in 1 hour at 37°C in a total reaction<br>volume of 50 μl. |
| <i>Lot</i> #:          | 0481709   |
| Assay Date:            | 09/2017   |
| Expiration Date:       | 9/2019  |
| Storage Temp:          | -20°C   |
| Storage Conditions:    | 50 mM NaCl, 10 mM Tris-HCl (pH 8.2), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.1% Tween-20, 200 $\mu$ g/ml BSA                                 |
| Specification Version: | PS-R0109S/L v1.0  |
| Effective Date:        | 14 Mar 2016   |

| Assay Name/Specification (minimum release criteria)   | Lot #0481709 |
|---|--------------|
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 5 units of FokI incubated for 4 hours at 37°C releases <0.5% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with FokI, $\sim$ 75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, $\sim$ 75% can be recut with FokI.  | Pass         |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

unon

Authorized by Derek Robinson 14 Mar 2016



Anthen The

Inspected by Anthony Francis 12 Sep 2017