

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | BanI   |
|------------------------|--|
| Catalog #:             | R0118S/L   |
| Concentration:         | 20,000 units/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Lot #:                 | 0271412  |
| Assay Date:            | 12/2014  |
| Expiration Date:       | 12/2016  |
| Storage Temp:          | -20 °C   |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA   |
| Specification Version: | PS-R0118S/L v1.0   |
| Effective Date:        | 17 Oct 2013  |

| Assay Name/Specification (minimum release criteria)  | Lot #0271412 |
|--|--------------|
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of BanI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.  | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with BanI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BanI.   | Pass         |
| <b>Non-Specific DNase Activity (16 hour)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of Lambda DNA and a minimum of 20 Units of BanI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme. | Pass         |
| <b>Protein Purity Assay (SDS-PAGE)</b> - BanI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.  | Pass         |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

M.W. Southworth

Authorized by Maurice Southworth 17 Oct 2013



Koeci Sakawaki

Inspected by Kerri Sakowski 31 Dec 2014