

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Sfil
Catalog Number:	R0123L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pXba in 1 hour at 50°C in a total reaction volume of 50 $\mu$ l.
Lot Number:	10013031
Expiration Date:	06/2020
Storage Temperature:	-20°C
Storage Conditions:	250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml BSA
Specification Version:	PS-R0123S/L v1.0

Sfil Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0123LVIAL	Sfil	10013014	Pass	
B7204SVIAL	CutSmart® Buffer	3091805	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0251805	Pass	

Assay Name/Specification	Lot # 10013031
Protein Purity Assay (SDS-PAGE) Sfil is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Sfil incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of Sfil incubated for 4 hours at 50°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pXba DNA with Sfil, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sfil.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10013031
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of pXba DNA and a minimum of	
100 units of Sfil incubated for 16 hours at 50°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis Production Scientist 22 Jun 2018

Much 1.

Michael Tonello Packaging Quality Control Inspector 25 Jun 2018

