

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sfil
Catalog Number: R0123L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba in 1 hour at 50°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10065029
Expiration Date: 02/2022
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0123S/L v1.0

Sfil Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0123LVIAL	Sfil	10065028	Pass	
B7204SVIAL	CutSmart® Buffer	10068803	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10064412	Pass	

Assay Name/Specification	Lot # 10065029
Protein Purity Assay (SDS-PAGE)	Pass
Sfil is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 100 units of Sfil incubated for 4 hours at 50°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 100 units of Sfil incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	
nours at 50°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pXba DNA with Sfil, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%	
can be recut with Sfil.	



R0123L / Lot: 10065029

Page 1 of 2



Assay Name/Specification	Lot # 10065029
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of	
100 units of Sfil incubated for 16 hours at 50°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

30 Jan 2020

Jay Minichiello

Packaging Quality Control Inspector

06 Mar 2020

