

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Sfil
Catalog Number:	R0123S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of pXba in 1 hour at 50°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10140927
Expiration Date:	03/2024
Storage Temperature:	-20°C
Storage Conditions:	250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml BSA
Specification Version:	PS-R0123S/L v1.0

Sfil Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0123SVIAL	Sfil	10140924	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10132772	Pass	
B6004SVIAL	rCutSmart™ Buffer	10138403	Pass	

Assay Name/Specification	Lot # 10140927
Protein Purity Assay (SDS-PAGE)	Pass
Sfil is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pXba DNA with Sfil, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sfil.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of	
100 units of Sfil incubated for 16 hours at 50°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 100 units of Sfil incubated for 4 hours at 50°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10140927
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³ H] E. coli DNA and a minimum of 100 units of Sfil incubated for 4	
hours at 50°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Panatha Thang

Penghua Zhang Production Scientist 28 Mar 2022

Michae

Michael Tonello Packaging Quality Control Inspector 28 Mar 2022

