

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	NlaIV
Catalog #:	R0126S/L
Concentration:	2,000 units/ml
-	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
<i>Lot</i> #:	0141308
Assay Date:	08/2013
Expiration Date:	08/2015
Storage Temp:	-20 °C
Storage Conditions:	200 mM KCl, 10 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0126S/L v1.0
Effective Date:	23 Jul 2013

Assay Name/Specification (minimum release criteria)	Lot #0141308
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 20 units of NlaIV incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of pBR322 DNA with NlaIV, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NlaIV.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of pBR322 DNA and a minimum of 20 Units of NlaIV incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

tunen

Authorized by Derek Robinson 23 Jul 2013



mo

Inspected by Penghua Zhang 21 Aug 2013