

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Pstl

Catalog Number: R0140M
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10056490
Expiration Date: 10/2021
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0140T/M v1.0

Pstl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0140MVIAL	PstI	10056489	Pass	
B7203SVIAL	NEBuffer™ 3.1	10053972	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10050274	Pass	

Assay Name/Specification	Lot # 10056490
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of Lambda DNA and a minimum of 100	Pass
units of PstI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE)	Pass
Pstl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Blue-White Screening (Terminal Integrity)	Pass
A sample of pUC19 vector linearized with a 10-fold excess of Pstl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in	
<1% white colonies.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and	
double-stranded [³ H] E. coli DNA and a minimum of 200 units of Pstl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
riours at 37°C releases <0.1% of the total radioactivity.	



R0140M / Lot: 10056490

Page 1 of 2



Assay Name/Specification	Lot # 10056490
Ligation and Recutting (Terminal Integrity)	Pass
After a 100-fold over-digestion of Lambda DNA with Pstl, >95% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Pstl.	

This product has been tested and shown to be in compliance with all specifications.

Production Scientist

26 Aug 2019

Michael Tonello

Packaging Quality Control Inspector

08 Oct 2019

