

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | PstI   |
|------------------------|--|
| Catalog #:             | R0140S/L   |
| Concentration:         | 20,000 units/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl. |
| <i>Lot</i> #:          | 0411510  |
| Assay Date:            | 10/2015  |
| Expiration Date:       | 10/2017  |
| Storage Temp:          | -20°C  |
| Storage Conditions:    | 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml<br>BSA                          |
| Specification Version: | PS-R0140S/L v1.0   |
| Effective Date:        | 08 May 2013  |

| Assay Name/Specification (minimum release criteria)  | Lot #0411510 |
|--|--------------|
| <b>Blue-White Screening (Terminal Integrity)</b> - A sample of pUC19 vector linearized with a 10-fold excess of PstI, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.  | Pass         |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of PstI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 100-fold over-digestion of Lambda DNA with PstI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PstI.  | Pass         |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of Lambda DNA and a minimum of 100 units of PstI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.     | Pass         |
| <b>Protein Purity Assay (SDS-PAGE)</b> - PstI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.  | Pass         |



R0140S/L Lot: 0411510 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otunen

Authorized by Derek Robinson 08 May 2013



Inspected by Jianying Luo 23 Oct 2015