

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Pstl
Catalog Number:	R0140T
Concentration:	100,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10140865
Expiration Date:	01/2024
Storage Temperature:	-20°C
Storage Conditions:	250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml BSA
Specification Version:	PS-R0140T/M v1.0

Pstl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0140TVIAL	Pstl	10134218	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10130600	Pass	
B6003SVIAL	NEBuffer™ r3.1	10132773	Pass	

Assay Name/Specification	Lot # 10140865
Protein Purity Assay (SDS-PAGE)	Pass
PstI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Blue-White Screening (Terminal Integrity)	Pass
A sample of pUC19 vector linearized with a 10-fold excess of Pstl, religated and	
transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of a mixture of single and	
double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of PstI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 100 units of PstI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10140865
Ligation and Recutting (Terminal Integrity)	Pass
After a 100-fold over-digestion of Lambda DNA with Pstl, >95% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Pstl.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 04 Mar 2022

Josh Hersey

Packaging Quality Control Inspector 04 Mar 2022

