

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pstl
Catalog Number: R0140T
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10183113
Expiration Date: 03/2025
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0140T/M v1.0

| Pstl Component List    |                              |            |                      |  |
|------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description        | Lot Number | Individual QC Result |  |
| R0140TVIAL             | PstI                         | 10183112   | Pass                 |  |
| B7024AVIAL             | Gel Loading Dye, Purple (6X) | 10182168   | Pass                 |  |
| B6003SVIAL             | NEBuffer™ r3.1               | 10168653   | Pass                 |  |

| Assay Name/Specification   | Lot # 10183113 |
|--|----------------|
| Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Pstl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in                                | Pass           |
| <1% white colonies.  Exonuclease Activity (Radioactivity Release)  | Pass           |
| A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 200 units of Pstl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.                      |                |
| Ligation and Recutting (Terminal Integrity) After a 100-fold over-digestion of Lambda DNA with Pstl, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Pstl. | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 100 units of Pstl incubated for 16 hours at 37°C results in a DNA pattern free of  | Pass           |



R0140T / Lot: 10183113

Page 1 of 2



| Assay Name/Specification   | Lot # 10183113 |
|--|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis.  |                |
| Protein Purity Assay (SDS-PAGE) Pstl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun \
Production Scientist

23 Mar 2023

Michael Tonello

Packaging Quality Control Inspector

27 Apr 2023



R0140T / Lot: 10183113

Page 2 of 2