

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	KpnI
Catalog #:	R0142M
Concentration:	50,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Lot #:	0551606
Assay Date:	06/2016
Expiration Date:	6/2018
Storage Temp:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0142M v1.0
Effective Date:	28 Mar 2014

Assay Name/Specification (minimum release criteria)	Lot #0551606
<b>Blue-White Screening (Terminal Integrity)</b> - A sample of pUC19 vector linearized with a 10-fold excess of KpnI, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> - A 50 $\mu$ l reaction in NEBuffer 1.1 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 30 Units of KpnI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in NEBuffer 1.1 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of KpnI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 20-fold over-digestion of pXba DNA with KpnI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with KpnI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 $\mu$ l reaction in NEBuffer 1.1 containing 1 $\mu$ g of pXba DNA and a minimum of 50 Units of KpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> - KpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass



R0142M Lot: 0551606 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otunen

Authorized by Derek Robinson 28 Mar 2014



Inspected by Jianying Luo 22 Jun 2016