

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Xbal
Catalog Number: R0145L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (dam-/HindIII digest) in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10104226
Expiration Date: 12/2022
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0145S/L v1.0

Xbal Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0145LVIAL	Xbal	10091826	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091457	Pass	
B6004SVIAL	rCutSmart™ Buffer	10103711	Pass	

Assay Name/Specification	Lot # 10104226
Protein Purity Assay (SDS-PAGE)	Pass
Xbal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII dam- DNA and a minimum of 200 Units of Xbal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 100 Units of Xbal incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 200 units of Xbal incubated for 4	Pass



R0145L / Lot: 10104226

Page 1 of 2

Assay Name/Specification	Lot # 10104226
hours at 37°C releases <0.1% of the total radioactivity.	
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Adenovirus-2 DNA with Xbal, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Xbal.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Xbal, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in 21% white colonies	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

13 Apr 2021

Michael Tonello

Packaging Quality Control Inspector

13 Apr 2021



R0145L / Lot: 10104226

Page 2 of 2