

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Mbol
Catalog Number: R0147L
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (dam-) in rCutSmart Buffer in 1 hour at 37°C in a

total reaction volume of 50 μl.

Packaging Lot Number: 10173510
Expiration Date: 12/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200

μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0147S/L v2.0

Mbol Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0147LVIAL	Mbol	10173506	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10168649	Pass	
B6004SVIAL	rCutSmart™ Buffer	10168651	Pass	

Assay Name/Specification	Lot # 10173510
Endonuclease Activity (Nicking) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of Mbol incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and a minimum of 50 units of Mbol incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1 µl of Mbol incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release)	Pass



R0147L / Lot: 10173510 Page 1 of 2

Assay Name/Specification	Lot # 10173510
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Mbol incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Mbol is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
Protein Purity Assay (SDS-PAGE) Mbol is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with Mbol, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Mbol.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun \
Production Scientist

08 Dec 2022

Michael Tonello

Packaging Quality Control Inspector

09 Dec 2022



R0147L / Lot: 10173510

Page 2 of 2