

New England Biolabs Certificate of Analysis

Product Name: TaqI-v2
Catalog Number: R0149T
Concentration: 100,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10079970
Expiration Date: 07/2022
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)
Specification Version: PS-R0149T/M v2.0

TaqI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0149TVIAL	TaqI-v2	10079969	Pass
B7204SVIAL	CutSmart® Buffer	10078757	Pass

Assay Name/Specification	Lot # 10079970
Protein Purity Assay (SDS-PAGE) TaqI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of TaqI-v2 is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of TaqI-v2 incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of TaqI-v2 incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

Assay Name/Specification	Lot # 10079970
<p>Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of TaqI-v2 incubated for 15 minutes at 65°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with TaqI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqI-v2.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
02 Oct 2020



Michael Tonello
Packaging Quality Control Inspector
02 Oct 2020