

www.neb.com info@neb.com



New England Biolabs Certificate of Analysis

Product Name: Avall
Catalog Number: R0153M
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total

reaction volume of 50 µl.

Packaging Lot Number: 10241247
Expiration Date: 02/2026
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200

μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0153T/M v2.0

Avail Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0153MVIAL	Avall	10229322	Pass	
B6004SVIAL	rCutSmart™ Buffer	10233337	Pass	

Assay Name/Specification	Lot # 10241247
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of AvaII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of Avall incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with AvaII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AvaII.	Pass
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of AvaII incubated for 16 hours at 37°C results in a DNA pattern free of	



R0153M / Lot: 10241247

Page 1 of 2



Assay Name/Specification	Lot # 10241247
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) AvaII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of AvaII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 25 Apr 2024 Michael Tonello

Packaging Quality Control Inspector

25 Apr 2024



R0153M / Lot: 10241247

Page 2 of 2