

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Sacl
Catalog Number: R0156L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10088069
Expiration Date: 09/2022
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0156S/L v1.0

SacI Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0156LVIAL	Sacl	10084607	Pass	
B7201SVIAL	NEBuffer™ 1.1	10090429	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084972	Pass	

Assay Name/Specification	Lot # 10088069
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Sacl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Sacl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of SacI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with Sacl, >95% of the DNA fragments can	Pass



R0156L / Lot: 10088069

Page 1 of 2

Assay Name/Specification	Lot # 10088069	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sacl.		
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of Lambda-HindIII DNA and a minimum of 60 units of Sacl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass	
Protein Purity Assay (SDS-PAGE) Sact is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection	Pass	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

24 Nov 2020

Josh Hersey

Packaging Quality Control Inspector

24 Nov 2020



R0156L / Lot: 10088069

Page 2 of 2