

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sacl
Catalog Number: R0156S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10084608
Expiration Date: 09/2022
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0156S/L v1.0

SacI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0156SVIAL	Sacl	10084606	Pass	
B7201SVIAL	NEBuffer™ 1.1	10065750	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084971	Pass	

Assay Name/Specification	Lot # 10084608
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of SacI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Sacl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of SacI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with Sacl, >95% of the DNA fragments can	Pass



R0156S / Lot: 10084608

Page 1 of 2

Assay Name/Specification	Lot # 10084608
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sacl.	
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of Lambda-HindIII DNA and a minimum of 60 units of Sacl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Sacl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pengha Zhang Production Scientist 27 Oct 2020 Michael Tonello

Packaging Quality Control Inspector

27 Oct 2020



R0156S / Lot: 10084608

Page 2 of 2