

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Sacli |
|------------------------|--|
| Catalog Number: | R0157L |
| Concentration: | 20,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μ g of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μ l. |
| Packaging Lot Number: | 10160453 |
| Expiration Date: | 08/2024 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R0157S/L v1.0 |

| SacII Component List | | | | |
|------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0157LVIAL | Sacll | 10160449 | Pass | |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10161525 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10158558 | Pass | |

| Assay Name/Specification | Lot # 10160453 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 60 Units of SacII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of SacII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 60 Units of SacII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with SacII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10160453 |
|--------------------------|----------------|
| can be recut with SacII. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist 09 Sep 2022

Erin Varney

Packaging Quality Control Inspector 09 Sep 2022

