

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: SacII
Catalog Number: R0157S
Concentration: 20,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10046226
Expiration Date: 06/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0157S/L v1.0

| SacII Component List   |                              |            |                      |  |
|------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description        | Lot Number | Individual QC Result |  |
| R0157SVIAL             | SacII                        | 10046227   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer             | 10043351   | Pass                 |  |
| B7024SVIAL             | Gel Loading Dye, Purple (6X) | 10043349   | Pass                 |  |

| Assay Name/Specification  | Lot # 10046226 |
|---|----------------|
| <b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with SacII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SacII.                                    | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 60 Units of SacII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 60 Units of SacII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.             | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of SacII incubated for  | Pass           |



R0157S / Lot: 10046226

Page 1 of 2



| Assay Name/Specification                                   | Lot # 10046226 |
|--|----------------|
| 4 hours at 37°C releases <0.1% of the total radioactivity. |                |

This product has been tested and shown to be in compliance with all specifications.

**Production Scientist** 

21 May 2019

Jay Minichiello

Packaging Quality Control Inspector

20 Jun 2019

