

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Bcll  |
|------------------------|---|
| Catalog Number:        | R0160S  |
| Concentration:         | 10,000 U/ml   |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg<br>of Lambda DNA (dam-) in 1 hour at 50°C in a total reaction volume of<br>50 μl. |
| Packaging Lot Number:  | 10147625  |
| Expiration Date:       | 04/2024   |
| Storage Temperature:   | -20°C   |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA   |
| Specification Version: | PS-R0160S/L v1.0  |

| Bcll Component List |                       |            |                      |  |
|---------------------|-----------------------|------------|----------------------|--|
| NEB Part Number     | Component Description | Lot Number | Individual QC Result |  |
| R0160SVIAL          | Bcll                  | 10147624   | Pass                 |  |
| B6003SVIAL          | NEBuffer™ r3.1        | 10146823   | Pass                 |  |

| Assay Name/Specification  | Lot # 10147625 |
|---|----------------|
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda dam- DNA and a minimum of 30 units of BcII incubated for 16 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BcII incubated for 4<br>hours at 50°C releases <0.1% of the total radioactivity.  | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 20-fold over-digestion of Lambda dam- DNA with BcII, >95% of the DNA<br>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated<br>fragments, >95% can be recut with BcII.                                  | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Penghua Zhang Production Scientist 27 Apr 2022

Erin Varney

Packaging Quality Control Inspector 27 Apr 2022

