

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Accl
Catalog Number: R0161L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10107872
Expiration Date: 05/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0161S/L v1.0

Accl Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0161LVIAL	Accl	10107870	Pass	
B6004SVIAL	rCutSmart™ Buffer	10107577	Pass	

Assay Name/Specification	Lot # 10107872
Protein Purity Assay (SDS-PAGE)	Pass
Accl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of Lambda DNA and a minimum of	
100 Units of Accl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
detectable hadreace degradation as determined by agained got electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with Accl, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Accl.	
Evenueleses Astivity (Redissativity Release)	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	Pass
double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Accl incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass



R0161L / Lot: 10107872

Page 1 of 2

Assay Name/Specification	Lot # 10107872
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled LITMUS28i DNA and a minimum of 100 Units of Accl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Accl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

09 Jun 2021

Michael Tonello

Packaging Quality Control Inspector

09 Jun 2021



R0161L / Lot: 10107872 Page 2 of 2