

## New England Biolabs Certificate of Analysis

**Product Name:** Accl  
**Catalog Number:** R0161L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10272954  
**Expiration Date:** 12/2026  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0161S/L/V v2.0

| Accl Component List |                       |            |                      |
|---------------------|-----------------------|------------|----------------------|
| NEB Part Number     | Component Description | Lot Number | Individual QC Result |
| R0161LVIAL          | Accl                  | 10268461   | Pass                 |
| B6004SVIAL          | rCutSmart™ Buffer     | 10265040   | Pass                 |

| Assay Name/Specification  | Lot # 10272954 |
|---|----------------|
| <b>Blue-White Screening (Terminal Integrity)</b><br>A sample of pUC19 vector linearized with a 10-fold excess of Accl, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.   | <b>Pass</b>    |
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled LITMUS28i DNA and a minimum of 100 units of Accl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.         | <b>Pass</b>    |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Accl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | <b>Pass</b>    |
| <b>Functional Testing (15 minute Digest)</b><br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of Accl incubated for 15 minutes at 37°C results in complete digestion as determined by  | <b>Pass</b>    |

| Assay Name/Specification  | Lot # 10272954 |
|---|----------------|
| agarose gel electrophoresis.  |                |
| <p><b>Ligation and Recutting (Terminal Integrity)</b><br/>After a 10-fold over-digestion of Lambda DNA with Accl, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with Accl.</p>   | <b>Pass</b>    |
| <p><b>Non-Specific DNase Activity (16 Hour)</b><br/>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Accl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>  | <b>Pass</b>    |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>Accl is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>   | <b>Pass</b>    |
| <p><b>qPCR DNA Contamination (E. coli Genomic)</b><br/>A minimum of 10 units of Accl is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> | <b>Pass</b>    |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Kyle Pepi  
Production Scientist  
19 Dec 2024



Michael Tonello  
Packaging Quality Control Inspector  
20 Dec 2024