

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | BstEII   |
|------------------------|--|
| Catalog #:             | R0162T/M   |
| Concentration:         | 50,000 units/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 60°C in a total reaction volume of 50 $\mu$ l. |
| Lot #:                 | 0851712  |
| Assay Date:            | 12/2017  |
| Expiration Date:       | 12/2019  |
| Storage Temp:          | -20°C  |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 $\mu$ g/ml BSA  |
| Specification Version: | <i>PS-R0162T/M v1.0</i>  |
| Effective Date:        | 23 Aug 2013  |

| Assay Name/Specification (minimum release criteria)   | Lot #0851712 |
|---|--------------|
| <b>Endonuclease Activity (Nicking)</b> - A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 30 units of BstEII incubated for 4 hours at 60°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.                  | Pass         |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of BstEII incubated for 4 hours at 60°C releases <0.1% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with BstEII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BstEII.  | Pass         |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 $\mu$ l reaction in NEBuffer 3.1 containing 1 $\mu$ g of Lambda DNA and a minimum of 50 Units of BstEII incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.     | Pass         |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otunen

Authorized by Derek Robinson 23 Aug 2013



en log

Inspected by Penghua Zhang 28 Dec 2017