

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: BstNI
Catalog Number: R0168S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 60°C in a total reaction volume of 50 μl.

Lot Number: 1001163.
Expiration Date: 06/2020
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0168S/L v1.0

| BstNI Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0168SVIAL | BstNI | 10011636 | Pass | |
| B7203SVIAL | NEBuffer™ 3.1 | 10021111 | Pass | |

| Assay Name/Specification | Lot # 10011635 |
|--|----------------|
| Exonuclease Activity (Radioactivity Release) | Pass |
| A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and | |
| double-stranded [3H] E. coli DNA and a minimum of 50 units of BstNI incubated for 4 | |
| hours at 60°C releases <0.1% of the total radioactivity. | |
| Ligation and Recutting (Terminal Integrity) | Pass |
| After a 2-fold over-digestion of Lambda DNA with BstNI, 95% can be recut with BstNI. | 1 335 |
| Non-Specific DNase Activity (16 Hour) | Pass |
| A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 30 | |
| Units of BstNI incubated for 16 hours at 60°C results in a DNA pattern free of | |
| detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.



R0168S / Lot: 10011635

Page 1 of 2

Stephani Onetto

Stephanie Cornelio Production Scientist 15 Jun 2018 Michael Tonello

Packaging Quality Control Inspector

08 Nov 2018