

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Sau3AI |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog Number: | R0169L |
| Concentration: | 5,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μ l. |
| Packaging Lot Number: | 10089314 |
| Expiration Date: | 11/2021 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R0169S/L v1.0 |

| Sau3AI Component List | | | | |
|-----------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0169LVIAL | Sau3AI | 10089315 | Pass | |
| B7201SVIAL | NEBuffer™ 1.1 | 10090429 | Pass | |

| Assay Name/Specification | Lot # 10089314 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 Units of Sau3AI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 15 units of Sau3AI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Sau3AI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sau3AI. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda DNA and a minimum of 25 Units of Sau3Al incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10089314 |
|------------------------------------------------------------------------------------------------------------------------------|----------------|
| Protein Purity Assay (SDS-PAGE) Sau3AI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 24 Nov 2020

Josh Hersey

Packaging Quality Control Inspector 24 Nov 2020

