

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Sau3Al
Catalog Number:	R0169L
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10122538
Expiration Date:	06/2022
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0169S/L v1.0

Sau3AI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0169LVIAL	Sau3AI	10111483	Pass	
B6001SVIAL	NEBuffer™ r1.1	10102943	Pass	

Assay Name/Specification	Lot # 10122538
Protein Purity Assay (SDS-PAGE) Sau3AI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 Units of Sau3Al incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 15 units of Sau3AI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda DNA and a minimum of 25 Units of Sau3Al incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10122538
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with Sau3AI, ~75% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Sau3AI.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Panatha Thang

Penghua Zhang Production Scientist 01 Oct 2021

Michae

Michael Tonello Packaging Quality Control Inspector 01 Oct 2021

