

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Sau3AI   |
|------------------------|--|
| Catalog Number:        | R0169S   |
| Concentration:         | 5,000 U/ml   |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Packaging Lot Number:  | 10210784   |
| Expiration Date:       | 09/2024  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0169S/L v1.0   |

| Sau3AI Component List |                       |            |                      |  |
|-----------------------|-----------------------|------------|----------------------|--|
| NEB Part Number       | Component Description | Lot Number | Individual QC Result |  |
| R0169SVIAL            | Sau3AI                | 10209141   | Pass                 |  |
| B6001SVIAL            | NEBuffer™ r1.1        | 10154035   | Pass                 |  |

| Assay Name/Specification   | Lot # 10210784 |
|--|----------------|
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a<br>minimum of 5 Units of Sau3Al incubated for 4 hours at 37°C results in <10%<br>conversion to the nicked form as determined by agarose gel electrophoresis.               | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 15 units of Sau3AI incubated for<br>4 hours at 37°C releases <0.1% of the total radioactivity.    | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 20-fold over-digestion of Lambda DNA with Sau3AI, ~75% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>>95% can be recut with Sau3AI.                                      | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda DNA and a minimum of 25<br>Units of Sau3Al incubated for 16 hours at 37°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification   | Lot # 10210784 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE)<br>Sau3AI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue<br>detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń

YunJie Sun Production Scientist 26 Sep 2023

Michae

Michael Tonello Packaging Quality Control Inspector 13 Oct 2023

