

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: AvrII

Catalog Number: R0174L

Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10119857
Expiration Date: 02/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R0174S/L v1.0

AvrII Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0174LVIAL	AvrII	10100185	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10114156	Pass	
B6004SVIAL	rCutSmart™ Buffer	10118974	Pass	

Assay Name/Specification	Lot # 10119857
Protein Purity Assay (SDS-PAGE)	Pass
AvrII is $\geq$ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in CutSmart <sup>τM</sup> Buffer containing 1 μg of Lambda HindIII DNA and a minimum of 50 Units of AvrII incubated for 16 hours at 37°C results in a DNA pattern	
free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 50-fold over-digestion of Lambda HindIII DNA with AvrII, >95% of the DNA	
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with AvrII.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	



R0174L / Lot: 10119857 Page 1 of 2

Assay Name/Specification	Lot # 10119857
a minimum of 50 Units of AvrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of AvrII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of AvrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang **Production Scientist** 

17 Sep 2021

Michael Tonello

Packaging Quality Control Inspector

17 Sep 2021



R0174L / Lot: 10119857 Page 2 of 2