

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Avrll
Catalog Number:	R0174S
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10054486
Expiration Date:	05/2021
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0174S/L v1.0

AvrII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0174SVIAL	Avrll	10044760	Pass	
B7204SVIAL	CutSmart® Buffer	10053981	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10047936	Pass	

Assay Name/Specification	Lot # 10054486
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 Units of AvrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of AvrII incubated for 4 hours at 37⁰C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 50-fold over-digestion of Lambda HindIII DNA with AvrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AvrII.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10054486
minimum of 50 Units of AvrII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) AvrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of AvrII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang

Penghua Zhang Production Scientist 14 May 2019

Michae MAI

Michael Tonello Packaging Quality Control Inspector 24 Sep 2019

