

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: AvrII

Catalog Number: R0174S

Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10081930 Expiration Date: 04/2022 Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0174S/L v1.0

AvrII Component List			
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result
R0174SVIAL	AvrII	10074012	Pass
B7204SVIAL	CutSmart® Buffer	10078752	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10082183	Pass

Assay Name/Specification	Lot # 10081930
Blue-White Screening (Terminal Integrity)	Pass
A sample of Litmus28i vector linearized with a 10-fold excess of AvrII, religated	
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
results in <1% write colonies.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 50 Units of AvrII incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 50 units of AvrII incubated for 4	
nours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 50-fold over-digestion of Lambda HindIII DNA with AvrII, >95% of the DNA	



R0174S / Lot: 10081930

Page 1 of 2

Assay Name/Specification	Lot # 10081930
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AvrII.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a minimum of 50 Units of AvrII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE)  AvrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist 03 Sep 2020

detection.

Michael Tonello

Packaging Quality Control Inspector

03 Sep 2020



R0174S / Lot: 10081930

Page 2 of 2