

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: AvrII

Catalog Number: R0174S

Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C

in a total reaction volume of 50 μl.

Packaging Lot Number: 10160785
Expiration Date: 06/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

500 μg/ml rAlbumin (pH 7.4 @25°C)

Specification Version: PS-R0174S/L/V v2.0

Avril Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0174SVIAL	AvrII	10152550	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10156431	Pass	
B6004SVIAL	rCutSmart™ Buffer	10156434	Pass	

Assay Name/Specification	Lot # 10160785
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of AvrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in rCutSmart <sup>TM</sup> Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of AvrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda-HindIII DNA and 1 μl of AvrII incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda-HindIII DNA and a	Pass



R0174S / Lot: 10160785

Page 1 of 2

Assay Name/Specification	Lot # 10160785
minimum of 50 units of AvrII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of AvrII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
Ligation and Recutting (Terminal Integrity) After a 50-fold over-digestion of Lambda HindIII DNA with AvrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AvrII.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of AvrII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Protein Purity Assay (SDS-PAGE) AvrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

07 Sep 2022

Erin Varney

Packaging Quality Control Inspector

07 Sep 2022



R0174S / Lot: 10160785

Page 2 of 2