

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Dpnl
Catalog Number: R0176L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10058274
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0176S/L v1.0

DpnI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0176LVIAL	DpnI	10053970	Pass	
B7204SVIAL	CutSmart® Buffer	10053983	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10053978	Pass	

Assay Name/Specification	Lot # 10058274
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnI.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass



R0176L / Lot: 10058274

Page 1 of 2

Assay Name/Specification	Lot # 10058274
Endenualege Activity (Nieking)	Page
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass
a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist

03 Sep 2019

Jay Minichiello

Packaging Quality Control Inspector

20 Nov 2019



R0176L / Lot: 10058274 Page 2 of 2