

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Dpnl
Catalog Number:	R0176L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10097366
Expiration Date:	01/2023
Storage Temperature:	-20°C
Storage Conditions:	400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0176S/L v1.0

DpnI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0176LVIAL	Dpnl	10097365	Pass	
B7204SVIAL	CutSmart® Buffer	10093122	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10091034	Pass	

Assay Name/Specification	Lot # 10097366	
Protein Purity Assay (SDS-PAGE)	Pass	
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.		
Non-Specific DNase Activity (16 Hour)	Pass	
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of		
100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.		
Exonuclease Activity (Radioactivity Release)	Pass	
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and		
double-stranded [³ H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.		
Endonuclease Activity (Nicking)	Pass	
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and		
a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10%		
onversion to the nicked form as determined by agarose gel electrophoresis.		





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10097366
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnI.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Peph Day

Penghua Zhang Production Scientist 12 Feb 2021

Josh Hersey

Packaging Quality Control Inspector 12 Feb 2021

