

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Dpnl
Catalog Number:	R0176L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10131735
Expiration Date:	12/2023
Storage Temperature:	-20°C
Storage Conditions:	400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0176S/L v1.0

DpnI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0176LVIAL	Dpnl	10131733	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10127723	Pass	
B6004SVIAL	rCutSmart™ Buffer	10130602	Pass	

Assay Name/Specification	Lot # 10131735
Protein Purity Assay (SDS-PAGE)	Pass
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnI.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of pBR322 DNA and a minimum of 100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10131735
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pepnolog

Penghua Zhang Production Scientist 05 Jan 2022

Mid

Michael Tonello Packaging Quality Control Inspector 05 Jan 2022

