

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Dpnl
Catalog Number: R0176L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10150217
Expiration Date: 02/2024
Storage Temperature: -20°C

Storage Conditions: 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0176S/L v1.0

DpnI Component List				
<b>NEB Part Number</b>	<b>Component Description</b>	Lot Number	Individual QC Result	
R0176LVIAL	DpnI	10137314	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10149690	Pass	
B6004SVIAL	rCutSmart™ Buffer	10146822	Pass	

Assay Name/Specification	Lot # 10150217
Protein Purity Assay (SDS-PAGE)	Pass
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Dpnl.	



R0176L / Lot: 10150217

Page 1 of 2

Assay Name/Specification	Lot # 10150217
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

16 May 2022

Erin Varney

Packaging Quality Control Inspector

16 May 2022



R0176L / Lot: 10150217

Page 2 of 2