

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Dpnl
Catalog Number: R0176S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10074286 Expiration Date: 08/2021 Storage Temperature: -20°C

Storage Conditions: 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0176S/L v1.0

DpnI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0176SVIAL	DpnI	10053971	Pass	
B7204SVIAL	CutSmart® Buffer	10074633	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10064412	Pass	

Assay Name/Specification	Lot # 10074286
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE)	Pass
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of	
100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	



R0176S / Lot: 10074286

Page 1 of 2



Assay Name/Specification	Lot # 10074286
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnI.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang

Production Scientist

12 Jun 2020

Michael Tonello

Packaging Quality Control Inspector

12 Jun 2020



R0176S / Lot: 10074286

Page 2 of 2