

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Dpnl
Catalog Number: R0176S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10077601 Expiration Date: 06/2022 Storage Temperature: -20°C

Storage Conditions: 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0176S/L v1.0

DpnI Component List				
<b>NEB Part Number</b>	<b>Component Description</b>	Lot Number	Individual QC Result	
R0176SVIAL	DpnI	10076066	Pass	
B7204SVIAL	CutSmart® Buffer	10075570	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10075964	Pass	

Assay Name/Specification	Lot # 10077601
Protein Purity Assay (SDS-PAGE)	Pass
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of	
100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
actionable hadioade degradation de determined by againede get electrophicionel	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with DpnI.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [ ³H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	



R0176S / Lot: 10077601

Page 1 of 2

Assay Name/Specification	Lot # 10077601
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang

**Production Scientist** 

16 Jul 2020

Michael Tonello

Packaging Quality Control Inspector

16 Jul 2020



R0176S / Lot: 10077601 Page 2 of 2