

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Dpnl
Catalog Number: R0176S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA (dam methylated) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10088440
Expiration Date: 08/2022
Storage Temperature: -20°C

Storage Conditions: 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0176S/L v1.0

DpnI Component List				
<b>NEB Part Number</b>	<b>Component Description</b>	Lot Number	Individual QC Result	
R0176SVIAL	DpnI	10082106	Pass	
B7204SVIAL	CutSmart® Buffer	10085423	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084971	Pass	

Assay Name/Specification	Lot # 10088440
Protein Purity Assay (SDS-PAGE)	Pass
DpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of	
100 units of DpnI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of pBR322 DNA with DpnI, ~25% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with DpnI.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 200 units of DpnI incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	



R0176S / Lot: 10088440

Page 1 of 2

Assay Name/Specification	Lot # 10088440
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 20 units of DpnI incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

03 Nov 2020

Michael Tonello

Packaging Quality Control Inspector

03 Nov 2020

