

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Xmal
Catalog Number:	R0180M
Concentration:	50,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g adenovirus-2 in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10107187
Expiration Date:	04/2023
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0180M v1.0

Xmal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0180MVIAL	Xmal	10107188	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089405	Pass	
B6004SVIAL	rCutSmart™ Buffer	10103711	Pass	

Assay Name/Specification	Lot # 10107187
Protein Purity Assay (SDS-PAGE) Xmal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of pXba DNA and a minimum of 10 Units of Xmal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Xmal incubated for 4 hours at 37 ^o C releases <0.2% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 Units of Xmal incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10107187
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pXba DNA with Xmal, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%	
can be recut with Xmal.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 30 Apr 2021

Michae

Michael Tonello Packaging Quality Control Inspector 30 Apr 2021

