

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Xmal
Catalog Number: R0180S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

pXba in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10121639
Expiration Date: 09/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0180S/L v1.0

Xmal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0180SVIAL	Xmal	10121638	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10121392	Pass	
B6004SVIAL	rCutSmart™ Buffer	10121395	Pass	

Assay Name/Specification	Lot # 10121639
Protein Purity Assay (SDS-PAGE)	Pass
Xmal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 10	
Units of Xmal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
dotottable hadioace degradation as determined by agained generation.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pXba DNA with Xmal, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Xmal.	
our se recut with Athai.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 50 units of Xmal incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	
Thousand at or o releases 50.270 of the total radioactivity.	



R0180S / Lot: 10121639

Page 1 of 2

Assay Name/Specification	Lot # 10121639
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 50 Units of Xmal incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

07 Oct 2021

Michael Tonello

Packaging Quality Control Inspector

07 Oct 2021



R0180S / Lot: 10121639

Page 2 of 2