

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Xmal
Catalog Number: R0180S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

pXba in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10140658
Expiration Date: 09/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0180S/L v1.0

| Xmal Component List | | | | |
|------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0180SVIAL | Xmal | 10121638 | Pass | |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10131976 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10136926 | Pass | |

| Assay Name/Specification | Lot # 10140658 |
|--|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Xmal incubated for 4 hours at 37°C releases <0.2% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pXba DNA with Xmal, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Xmal. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 10 Units of Xmal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Protein Purity Assay (SDS-PAGE) Xmal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | Pass |



R0180S / Lot: 10140658 Page 1 of 2

| Assay Name/Specification | Lot # 10140658 |
|---|----------------|
| Endonuclease Activity (Nicking) | Pass |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and | |
| a minimum of 50 Units of Xmal incubated for 4 hours at 37°C results in <10% | |
| conversion to the nicked form as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

22 Feb 2022

Michael Tonello

Packaging Quality Control Inspector

22 Feb 2022



R0180S / Lot: 10140658

Page 2 of 2