

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Stul

Catalog Number: R0187S

Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10151296 Expiration Date: 03/2024 Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0187S/L v1.0

Stul Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0187SVIAL	Stul	10145435	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10149690	Pass	
B6004SVIAL	rCutSmart™ Buffer	10150371	Pass	

Assay Name/Specification	Lot # 10151296
Protein Purity Assay (SDS-PAGE) Stul is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart <sup>™</sup> Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 300 units of Stul incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 10 Units of Stul incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus 28i vector linearized with a 10-fold excess of Stul, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass



R0187S / Lot: 10151296

Page 1 of 2

Assay Name/Specification	Lot # 10151296
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Stul, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Stul.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of Stul incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist 19 May 2022 Erin Varney

Packaging Quality Control Inspector

19 May 2022



R0187S / Lot: 10151296

Page 2 of 2