

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EcoRV
Catalog Number: R0195T
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10032365Expiration Date:12/2020Storage Temperature:-20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0195T/M v1.0

EcoRV Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0195TVIAL	EcoRV	10032367	Pass	
B7203SVIAL	NEBuffer™ 3.1	10021113	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021135	Pass	

Assay Name/Specification	Lot # 10032365
Protein Purity Assay (SDS-PAGE) EcoRV is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	_
A sample of LITMUS28i vector linearized with a 10-fold excess of EcoRV, religated	Pass
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a	Pass
minimum of 20 units of EcoRV incubated for 4 hours at 37°C results in <50% conversion to the nicked form as determined by agarose gel electrophoresis.	
	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and	Гаээ
double-stranded [³H] E. coli DNA and a minimum of 100 units of EcoRV incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R0195T / Lot: 10032365

Page 1 of 2

Assay Name/Specification	Lot # 10032365
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with EcoRV, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRV.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 20 units of EcoRV incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso **Production Scientist**

NEW ENGLAND

26 Nov 2018

Michael Tonello

Packaging Quality Control Inspector

14 Jan 2019



R0195T / Lot: 10032365

Page 2 of 2