

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: RsrII
Catalog Number: R0501L
Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10137472
Expiration Date: 07/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.15% Triton® X-100, 200 μg/ml BSA (pH 7.4 @ 25°C)

Specification Version: PS-R0501S/L v2.0

Rsrll Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0501LVIAL	Rsrll	10137471	Pass	
B6004SVIAL	rCutSmart™ Buffer	10154051	Pass	

Assay Name/Specification	Lot # 10137472
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 50 units of RsrII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protein Purity Assay (SDS-PAGE) RsrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



R0501L / Lot: 10137472

Page 1 of 2

Assay Name/Specification	Lot # 10137472
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with RsrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with RsrII.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

29 Jul 2022

Erin Varney

Packaging Quality Control Inspector

29 Jul 2022



R0501L / Lot: 10137472

Page 2 of 2