

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: RsrII
Catalog Number: R0501L
Concentration: 5,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10181039
Expiration Date: 03/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.15% Triton® X-100, 200  $\mu$ g/ml BSA (pH 7.4 @ 25°C)

Specification Version: PS-R0501S/L v2.0

RsrII Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0501LVIAL	Rsrll	10181038	Pass	
B6004SVIAL	rCutSmart™ Buffer	10181134	Pass	

Assay Name/Specification	Lot # 10181039
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass
a minimum of 50 units of RsrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with RsrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with RsrII.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 50 units of RsrII incubated for 16 hours at 37°C results in a DNA pattern free of	Pass



R0501L / Lot: 10181039

Page 1 of 2



Assay Name/Specification	Lot # 10181039
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) RsrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist 28 Feb 2023

Michael Tonello

Packaging Quality Control Inspector

22 May 2023



R0501L / Lot: 10181039

Page 2 of 2